

# Differentiable Manifolds

## §18. Differentiable $k$ -Forms

Sichuan University, Fall 2022

## Reminder (see Section 3)

Let  $V$  be a vector space (over  $\mathbb{R}$ ). Set  $n = \dim V$ .

- A  $k$ -covector on  $V$  is an alternating  $k$ -linear map  $f : V^k \rightarrow \mathbb{R}$ ,

$$f(v_{\sigma(1)}, \dots, v_{\sigma(k)}) = (\operatorname{sgn} \sigma) f(v_1, \dots, v_n) \quad \forall \sigma \in S_k.$$

- We denote by  $A_k(V)$  the space of  $k$ -covectors on  $V$ .
- We have

$$A_0(V) = \mathbb{R}, \quad A_1(V) = V^*, \quad A_k(V) = \{0\}, \quad k \geq n+1.$$

## Reminder (Wedge product; see Section 3)

- If  $f \in A_k(V)$  and  $g \in A_\ell(V)$ , the *wedge product*  $f \wedge g$  is the  $(k + \ell)$ -covector in  $A_{k+\ell}(V)$  defined by

$$(f \wedge g)(v_1, \dots, v_{k+\ell}) = \frac{1}{k!\ell!} \sum_{\sigma \in S_{k+\ell}} \text{sgn}(\sigma) f(v_{\sigma(1)}, \dots, v_{\sigma(k)}) g(v_{\sigma(k+1)}, \dots, v_{\sigma(k+\ell)}).$$

- The wedge product  $\wedge : A_k(V) \times A_\ell(V) \rightarrow A_{k+\ell}(V)$  is a bilinear map which is anti-commutative and associative, i.e.,

$$f \wedge g = (-1)^{k\ell} g \wedge f, \quad f \wedge f = 0 \quad (k \text{ odd}), \\ (f \wedge g) \wedge h = f \wedge (g \wedge h).$$

## Reminder (Wedge products of 1-covectors; see Section 3)

- If  $\alpha^1, \dots, \alpha^k$  are 1-covectors, then

$$(\alpha^1 \wedge \dots \wedge \alpha^k)(v_1, \dots, v_k) = \det [\alpha^i(v_j)], \quad v_i \in V.$$

- Let  $\beta^1, \dots, \beta^k$  be  $k$ -covectors such that

$$\beta^i = \sum_j a_j^i \alpha^j, \quad \text{for some matrix } A = [a_j^i] \in \mathbb{R}^{k \times k}.$$

Then

$$\beta^1 \wedge \dots \wedge \beta^k = (\det A) \alpha^1 \wedge \dots \wedge \alpha^k.$$

# Differential Forms

## Definition

$\mathcal{I}_{k,n}$  is the set of ascending multi-indices  $I = (i_1, \dots, i_k)$  such that  $1 \leq i_1 < \dots < i_k \leq n$ .

## Reminder (Bases of $k$ -covectors; see Section 3)

Let  $e_1, \dots, e_n$  be a basis of  $V$  and let  $\alpha^1, \dots, \alpha^n$  be the dual basis of  $V^* = A_1(V)$ . For  $I = (i_1, \dots, i_k) \in \mathcal{I}_{k,n}$  set

$$\alpha^I = \alpha^{i_1} \wedge \dots \wedge \alpha^{i_k}.$$

- If  $J = (j_1, \dots, j_k) \in \mathcal{I}_{k,n}$  and  $e_J = (e_{j_1}, \dots, e_{j_k})$ , then

$$\alpha^I(e_J) = \delta^I_J.$$

- The  $k$ -covectors  $\alpha^I$ ,  $I \in \mathcal{I}_{k,n}$ , form a basis of  $A_k(V)$ .
- In particular  $\dim A_k(V) = \binom{n}{k}$  for  $k \leq n$ .

# Differential Forms

## Facts

- Any linear map  $F : V \rightarrow W$  gives rise to a linear map  $F^* : A_k(W) \rightarrow A_k(V)$  defined by

$$F^*g(v_1, \dots, v_k) = g(Fv_1, \dots, Fv_k), \quad g \in A_k(W), \quad v_i \in V.$$

- If  $F : V \rightarrow W$  and  $G : W \rightarrow Z$  are linear maps, then

$$(G \circ F)^* = F^* \circ G^*.$$

## Consequence

The construction  $V \rightarrow A_k(V)$  is a (contravariant) functor from the category  $\mathbf{Vect}_{\mathbb{R}}$  to itself.

## Remark

- There is another construction  $V \rightarrow \Lambda^k(V)$  called  *$k$ -th exterior power*.
- This is a covariant functor on  $\mathbf{Vect}_{\mathbb{R}}$ .
- We have  $A_k(V) = \Lambda^k(V^*)$ , so the space of  $k$ -covectors is often denoted  $\Lambda^k(V^*)$ .

# Differential Forms

## Definition (Differential $k$ -forms)

Let  $M$  be a smooth manifold.

- The space  $A_k(T_p M)$  is denoted  $\Lambda^k(T_p^* M)$ .
- An element of  $\Lambda^k(T_p^* M)$  is called a  $k$ -covector at  $p$ .
- A *differential  $k$ -form* (or a  *$k$ -covector field*) is the assignment for each  $p \in M$  of a  $k$ -covector  $\omega \in \Lambda^k(T_p^* M)$ .

## Remarks

- 1 Differential  $k$ -forms are also called *differential forms of degree  $k$* , or simply  *$k$ -forms*.
- 2 A differential form of degree  $k = \dim M$  is also called a *top form*.



# Differential Forms

## Definition

If  $\omega$  is a differential  $k$ -form and  $X_1, \dots, X_k$  are vector fields on  $M$ , we denote by  $\omega(X_1, \dots, X_k)$  the function on  $M$  defined by

$$\omega(X_1, \dots, X_k)(p) = \omega_p((X_1)_p, \dots, (X_k)_p), \quad p \in M.$$

## Proposition (Proposition 18.1)

Let  $\omega$  be a differential  $k$ -form. For any vector fields  $X_1, \dots, X_k$  and function  $h$  on  $M$ , we have

$$\omega(X_1, \dots, hX_i, \dots, X_k) = h\omega(X_1, \dots, X_k).$$

## Example

Let  $(U, x^1, \dots, x^n)$  be a chart for  $M$ .

- If  $p \in U$ , then  $\{\partial/\partial x^1|_p, \dots, \partial/\partial x^n|_p\}$  is a basis of  $T_p M$ .
- The dual basis of  $T_p^* M$  is  $\{(dx^1)_p, \dots, (dx^n)_p\}$ .
- For  $I = (i_1, \dots, i_k) \in \mathcal{I}_{k,n}$  let  $dx^I$  be the  $k$ -form defined by

$$(dx^I)_p = (dx^{i_1})_p \wedge \dots \wedge (dx^{i_k})_p, \quad p \in U.$$

By the results of Section 3 (see slide 5)  $\{(dx^I)_p; I \in \mathcal{I}_{k,n}\}$  is a basis of  $\Lambda^k(T_p^* M)$  for every  $p \in U$ .

# Local Expression for a $k$ -Form

## Facts

- Let  $p \in U$ . As  $\{(dx^I)_p; I \in \mathcal{I}_{k,n}\}$  is a basis of  $\Lambda^k(T_p^*M)$ , every  $k$ -covector  $\omega_p \in \Lambda^k(T_p^*M)$  can be uniquely written as

$$\omega_p = \sum_{I \in \mathcal{I}_{k,n}} a_I (dx^I)_p, \quad a_I \in \mathbb{R}.$$

- Set  $\partial_i = \partial/\partial x^i$  and for  $I = (i_1, \dots, i_k) \in \mathcal{I}_{k,n}$  set  $\partial_I = (\partial_{i_1}, \dots, \partial_{i_k})$ . By the results of Section 3 (see slide 5):

$$dx^I(\partial_J) = \delta_J^I.$$

It follows that if  $\omega_p = \sum_{I \in \mathcal{I}_{k,n}} a_I (dx^I)_p$ , then  $a_I = \omega_p(\partial_I)$ .

- In particular, every  $k$ -form  $\omega$  on  $U$  can be uniquely written as

$$\omega = \sum_{I \in \mathcal{I}_{k,n}} a_I dx^I \quad \text{with } a_I = \omega(\partial_I).$$

# Local Expression for a $k$ -Form

## Proposition (Proposition 18.3)

Suppose that  $(U, x^1, \dots, x^n)$  is a chart for  $M$ , and let  $f^1, \dots, f^k$  be smooth functions on  $U$ . Then

$$df^1 \wedge \dots \wedge df^k = \sum_I \frac{\partial(f^1, \dots, f^k)}{\partial(x^{i_1}, \dots, x^{i_k})} dx^I.$$

## Remark

In fact, in the same way as in Section 3 (see slide 4), we have

$$\begin{aligned} (df^1 \wedge \dots \wedge df^k)(\partial_{i_1}, \dots, \partial_{i_k}) &= \det [df^i(\partial_{i_j})] = \det [\partial f^i / \partial x^{i_j}] \\ &= \frac{\partial(f^1, \dots, f^k)}{\partial(x^{i_1}, \dots, x^{i_k})}. \end{aligned}$$

# Local Expression for a $k$ -Form

## Example

Let  $(V, y^1, \dots, y^n)$  be another chart. Then on  $U \cap V$  we have

$$dy^J = \sum_I \frac{\partial(y^{j_1}, \dots, y^{j_k})}{\partial(x^{i_1}, \dots, x^{i_k})} dx^I.$$

## Corollary (Corollary 18.4)

Suppose that  $(U, x^1, \dots, x^n)$  is a chart for  $M$ , and let  $f, f^1, \dots, f^n$  be smooth functions on  $U$ . Then

$$df = \sum_i \frac{\partial f}{\partial x^i} dx^i,$$
$$df^1 \wedge \dots \wedge df^n = \frac{\partial(f^1, \dots, f^n)}{\partial(x^1, \dots, x^n)} dx^1 \wedge \dots \wedge dx^n.$$

## Definition

- The  $k$ -th exterior power of the cotangent bundle is

$$\Lambda^k(T^*M) = \bigsqcup_{p \in M} \Lambda^k(T_p^*M) = \left\{ (p, \omega); \ p \in M, \ \omega \in \Lambda^k(T_p^*M) \right\}.$$

- The canonical map  $\pi : \Lambda^k(T^*M) \rightarrow M$  is given by

$$\pi(p, \omega) = p, \quad p \in M, \quad \omega \in \Lambda^k(T_p^*M).$$

# The Bundle Point of View

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ . Set  $V = \phi(U)$ .

- Every  $k$ -covector  $\omega_p \in \Lambda^k(T_p^*M)$  can be uniquely written as

$$\omega_p = \sum_I a_I (dx^I)_p, \quad \text{with } a^I = \omega_p(\partial_I).$$

- We thus get a natural bijection  $\tilde{\phi} : \Lambda^k(T^*U) \rightarrow V \times \mathbb{R}^{\binom{n}{k}}$  such that, for all  $p \in M$  and  $\omega \in \Lambda^k(T_p^*M)$ , we have

$$\tilde{\phi}(p, \omega) = ((x^i(p)), (\omega(\partial_I))).$$

## Remark

In the same way as with the constructions of the tangent bundle  $TM$  and the cotangent bundle  $T^*M$ , the maps  $\tilde{\phi}$  allow us to define a topology and a smooth structure on  $\Lambda^k(T^*M)$ .

# The Bundle Point of View

## Definition

Let  $(U, \phi)$  be a chart for  $M$  and set  $V = \phi(U)$ . We endow  $\Lambda^k(T^*U)$  with the topology such that

$$W \subset \Lambda^k(T^*U) \text{ is open} \iff \tilde{\phi}(W) \text{ is open in } V \times \mathbb{R}^{\binom{n}{k}}.$$

## Proposition

Let  $\{(U_\alpha, \phi_\alpha)\}$  be the maximal atlas of  $M$ .

- Define

$$\mathcal{B} = \bigcup_{\alpha} \left\{ W; W \text{ is an open in } \Lambda^k(T^*U_\alpha) \right\}.$$

Then  $\mathcal{B}$  is the basis for a unique topology on  $\Lambda^k(T^*M)$ .

- The collection  $\{(T^*U_\alpha, \tilde{\phi}_\alpha)\}$  is a  $C^\infty$  atlas on  $\Lambda^k(T^*M)$ , and hence  $\Lambda^k(T^*M)$  is a smooth manifold.
- $\Lambda^k(T^*M) \xrightarrow{\pi} M$  is a smooth vector bundle over  $M$ .



# Smooth $k$ -Forms

## Remark

A  $k$ -form on  $M$  is a section of the exterior power  $\Lambda^k(T^*M)$ .

## Definition

- We say that  $k$ -form is  $C^\infty$  when it is  $C^\infty$  as a section of  $\Lambda^k(T^*M)$ .
- We denote by  $\Omega^k(M)$  the space of smooth  $k$ -forms on  $M$ .

## Remarks

- ① In other words,  $\Omega^k(M)$  is the space of smooth sections of  $T^*M$ . In particular, this is a module over the ring  $C^\infty(M)$ .
- ② As  $\Lambda^0(T_p^*M) = \mathbb{R}$ , a 0-form is just a map from  $M$  to  $\mathbb{R}$ . Thus, a smooth 0-form is just a smooth function on  $M$ , i.e.,  $\Omega^0(M) = C^\infty(M)$ .

# Smooth $k$ -Forms

## Example

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ . Set  $V = \phi(U)$ .

- It can be shown that each  $k$ -form  $dx^I$ ,  $I \in \mathcal{I}_{k,n}$  is smooth.
- Thus,  $\{dx^I; I \in \mathcal{I}_{n,k}\}$  is a smooth frame of  $\Lambda^k(T^*M)$  over  $U$ .

## Reminder (Proposition 12.2)

Let  $\{s_1, \dots, s_r\}$  be a  $C^\infty$  frame of a vector bundle  $E$  over  $U$ . A section  $s = \sum c^i s_i$  of  $E$  over  $U$  is smooth if and only if  $c^1, \dots, c^r$  are smooth functions on  $U$ .

We immediately obtain:

## Lemma (Lemma 18.6)

Let  $(U, x^1, \dots, x^n)$  be a chart for  $M$ . A  $k$ -form  $\omega = \sum a_I dx^I$  on  $U$  is smooth if and only if the coefficients  $a_I$  are  $C^\infty$  functions on  $U$ .

# Smooth $k$ -Forms

In the same way as with vector fields and 1-forms by using the previous lemma we obtain:

## Proposition (Proposition 18.7; 1st part)

Let  $\omega$  be a  $k$ -form on  $M$ . Then TFAE:

- ①  $\omega$  is a smooth  $k$ -form.
- ②  $M$  has an atlas such that, for every chart  $(U, x^1, \dots, x^n)$  of this atlas, we may write  $\omega = \sum a_I dx^I$  on  $U$  with  $a^I \in C^\infty(U)$ .
- ③ For every chart  $(U, x^1, \dots, x^n)$  of  $M$ , we may write  $\omega = \sum a_I dx^I$  on  $U$  with  $a^I \in C^\infty(U)$ .

# Smooth $k$ -Forms

## Proposition (Proposition 18.7; 2nd part)

Let  $\omega$  be a  $k$ -form on  $M$ . Then TFAE:

- ①  $\omega$  is a smooth  $k$ -form.
- ② For any smooth vector fields  $X_1, \dots, X_k$  on  $M$ , the function  $\omega(X_1, \dots, X_k)$  is smooth on  $M$ .

## Proposition (Proposition 18.8)

Let  $\tau$  be a smooth  $k$ -form defined on a neighborhood of  $p$ . Then there exists a smooth  $k$ -form  $\tilde{\tau}$  on  $M$  which agrees with  $\tau$  near  $p$ .

# Pullback of $k$ -Forms

Reminder (see slide 6)

Any linear  $F : V \rightarrow W$  between vector spaces gives rise to a linear map  $F^* : A_k(W) \rightarrow A_k(V)$  defined by

$$F^*g(v_1, \dots, v_k) = g(Fv_1, \dots, Fv_k), \quad g \in A_k(W), \quad v_i \in V.$$

Definition (Pullback of a  $k$ -form)

Let  $F : N \rightarrow M$  be a smooth map. If  $\omega$  is a  $k$ -form on  $M$ , then its pullback  $F^*\omega$  is the  $k$ -form on  $N$  defined by

$$(F^*\omega)_p = (F_{*,p})^* \omega_{F(p)}, \quad p \in N.$$

That is,

$$(F^*\omega)_p(v_1, \dots, v_k) = \omega_p(F_{*,p}v_1, \dots, F_{*,p}v_k), \quad v_i \in T_pM.$$

# Pullback of $k$ -Forms

## Proposition (Proposition 18.9)

Let  $F : N \rightarrow M$  be a smooth map. If  $\omega$  and  $\tau$  are  $k$ -forms on  $M$  and  $a$  is a constant, then

$$\begin{aligned}F^*(\omega + \tau) &= F^*\omega + F^*\tau, \\F^*(a\omega) &= aF^*\omega.\end{aligned}$$

## Remark

We will see later that if  $\omega$  is a smooth  $k$ -form, then its pullback  $F^*\omega$  is a smooth  $k$ -form as well (see slide 29).

# The Wedge Product

## Definition

If  $\omega$  is a  $k$ -form and  $\tau$  is a  $\ell$ -form on  $M$ , then their *wedge product*  $\omega \wedge \tau$  is the  $(k + \ell)$ -form on  $M$  defined by

$$(\omega \wedge \tau)_p = \omega_p \wedge \tau_p \in \Lambda^{k+\ell}(T_p^*M), \quad p \in M.$$

## Proposition (Proposition 18.10)

If  $\omega$  and  $\tau$  are smooth forms on  $M$ , then  $\omega \wedge \tau$  is smooth on  $M$ .

## Corollary

The wedge product induces an anti-commutative associative bilinear map,

$$\wedge : \Omega^k(M) \times \Omega^\ell(M) \longrightarrow \Omega^{k+\ell}(M).$$

# The Wedge Product

## Reminder (Graded Algebras)

- ① An algebra  $A$  over a field  $\mathbb{K}$  is called *graded* when it can be decomposed as

$$A = \bigoplus_{k=0}^{\infty} A^k,$$

where the  $A^k$  are subspaces such that the multiplication maps  $A^k \times A^\ell$  to  $A^{k+\ell}$ .

- ② We say that  $A$  is *anticommutative* (or *graded commutative*) when

$$ba = (-1)^{k\ell} ab \quad \text{for all } a \in A^k \text{ and } b \in A^\ell.$$



# The Wedge Product

## Proposition

Set  $n = \dim M$ . We Define

$$\Omega^*(M) = \bigoplus_{k=0}^n \Omega^k(M).$$

Then  $\Omega^*(M)$  is anticommutative graded algebra under the wedge product.

## Remark

$\Omega^*(M)$  is called the exterior algebra of differential forms on  $M$ .

# Wedge Product and Pullback

## Proposition (Proposition 18.11)

Let  $F : N \rightarrow M$  be a smooth map. If  $\omega$  and  $\tau$  are differential forms on  $M$ , then

$$F^*(\omega \wedge \tau) = (F^*\omega) \wedge (F^*\tau).$$

This result is used to prove:

## Lemma (Local expression for pullback)

Suppose that  $F : N \rightarrow M$  is a smooth map. Let  $(U, x^1, \dots, x^n)$  be a chart for  $N$  and  $(V, y^1, \dots, y^m)$  a chart for  $M$  such that  $U \subset F^{-1}(V)$ . Set  $F^j = y^j \circ F$ . For any  $k$ -form  $\omega = \sum b_J dy^J$  on  $V$ , we have

$$F^*\omega = \sum_{I,J} (b_J \circ F) \frac{\partial(F^{j_1}, \dots, F^{j_k})}{\partial(x^{i_1}, \dots, x^{i_k})} dx^I \quad \text{on } U.$$

# Wedge Product and Pullback

Proof.

- Thanks to Proposition 18.9, on  $F^{-1}(V)$  we have

$$F^*\omega = F^*\left(\sum_J b_J y^J\right) = \sum_J F^*b_J F^*(dy^J) = \sum_J (b_J \circ F) F^*(dy^J).$$

- It remains to determine  $F^*(dy^J)$ . By Proposition 18.11,  
$$F^*(dy^J) = F^*(dy^{j_1} \wedge \dots \wedge dy^{j_k}) = (F^*dy^{j_1}) \wedge \dots \wedge (F^*dy^{j_k}).$$

- By Proposition 17.10 pullback commutes with the differential:

$$(F^*dy^{j_\ell}) = d(F^*y^{j_\ell}) = d(y^{j_\ell} \circ F) = dF^{j_\ell}.$$

- Thus, on  $U$  we have

$$F^*(dy^J) = dF^{j_1} \wedge \dots \wedge dF^{j_k} = \sum_I \frac{\partial(F^{j_1}, \dots, F^{j_k})}{\partial(x^{i_1}, \dots, x^{i_k})} dx^I.$$

This gives the result. □

# Wedge Product and Pullback

By combining the previous lemma with the characterization of smoothness of  $k$ -forms (Proposition 18.7) we obtain:

## Proposition (Proposition 19.7)

*Let  $F : N \rightarrow M$  be a smooth map. If  $\omega$  is a smooth  $k$ -form on  $M$ , then  $F^*\omega$  is a smooth form on  $N$ .*

## Remark

- In Tu's book the above result is proved in Section 19. The main step is to prove the previous lemma.
- However, Tu's proof uses Proposition 19.5 whose statement requires Proposition 19.7 in order to make sense.
- Therefore, Proposition 19.5 cannot be used to prove Proposition 19.7.
- Tu's arguments are fine if we use Proposition 17.10 instead of Proposition 19.5 (as it is done in the previous slide).

# Invariant Forms on a Lie Group

## Definition

Let  $G$  be a Lie group. A  $k$ -form  $\omega$  on  $G$  is *left-invariant* if

$$\ell_g^* \omega = \omega \quad \forall g \in G,$$

where  $\ell_g : G \rightarrow G$  is the left-multiplication by  $g$ .

## Remark

The left-invariance condition means that

$$(\ell_g)_{*,x}^* (\omega_{gx}) = \omega_x \quad \forall g, x \in G.$$

In particular, by substituting  $g$  for  $x$  and  $g^{-1}$  for  $g$  we get

$$\omega_g = (\ell_{g^{-1}})_{*,g}^* (\omega_e) \quad \forall g \in G.$$

Thus,  $\omega$  is uniquely determined by  $\omega_e$ .

# Invariant Forms on a Lie Group

## Remark

Any  $k$ -covector  $\omega \in \Lambda^k(T_e^*G)$  generates a left-invariant  $k$ -form  $\tilde{\omega}$  defined by

$$\tilde{\omega}_g = (\ell_{g^{-1}})_{*,g}^*(\omega), \quad g \in G.$$

## Proposition (Proposition 18.14)

*Every left-invariant  $k$ -form on  $G$  is smooth.*

## Consequence

Denote by  $\Omega^k(M)^G$  the space of left-invariant  $k$ -forms on  $G$ . Then we have a linear isomorphism,

$$\Omega^k(G)^G \longrightarrow \Lambda^k(T_e^*G), \quad \omega \longrightarrow \omega_e.$$

In particular, if  $n = \dim G$ , then  $\Omega^k(G)^G$  has dimension  $\binom{n}{k}$ .

# Differential Forms on $\mathbb{S}^1$

## Proposition (Problem 18.8)

Let  $F : N \rightarrow M$  be a surjective submersion.

- The pullback by  $F$  gives rise to an injective linear map  $F^* : \Omega^k(M) \rightarrow \Omega^k(N)$ .
- This allows us to identify  $\Omega^k(M)$  with a subspace of  $\Omega^k(N)$ .

## Definition

- A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a  $2\pi$ -periodic if  $f(t + 2\pi) = f(t)$ .
- A 1-form  $f(t)dt$  on  $\mathbb{R}$  is said to be  $2\pi$ -periodic if the function  $f(t)$  is  $2\pi$ -periodic.

## Proposition (Proposition 18.12)

Let  $h : \mathbb{R} \rightarrow \mathbb{S}^1$  be the map defined by

$$h(t) = (\cos t, \sin t).$$

Then:

- $h$  is a surjective submersion.
- For  $k = 0, 1$ , under the pullback map  $h^* : \Omega^k(\mathbb{S}^1) \rightarrow \Omega^k(\mathbb{R})$  the smooth  $k$ -forms on  $\mathbb{S}^1$  corresponds to smooth  $2\pi$ -periodic  $k$ -forms on  $\mathbb{R}$ .