

# Differentiable Forms in Algebraic Topology

## Review: The Tangent Bundle

Sichuan University, Fall 2024

# The Tangent Bundle as a Manifold

## Objective

Let  $M$  be smooth manifold of dimension  $n$ . We would like to bundle together all the tangent spaces  $T_p M$  so as to get a smooth manifold, called the *tangent bundle*.

# The Tangent Bundle as a Manifold

## Objective

Let  $M$  be smooth manifold of dimension  $n$ . We would like to bundle together all the tangent spaces  $T_p M$  so as to get a smooth manifold, called the *tangent bundle*.

## Definition

As a set, the *tangent bundle* of  $M$  is the disjoint union,

$$TM := \bigsqcup_{p \in M} T_p M = \{(p, v); p \in M, v \in T_p M\}.$$

# The Tangent Bundle as a Manifold

## Objective

Let  $M$  be smooth manifold of dimension  $n$ . We would like to bundle together all the tangent spaces  $T_p M$  so as to get a smooth manifold, called the *tangent bundle*.

## Definition

As a set, the *tangent bundle* of  $M$  is the disjoint union,

$$TM := \bigsqcup_{p \in M} T_p M = \{(p, v); p \in M, v \in T_p M\}.$$

## Remarks

- 1 For  $p \in M$  we identify the subset  $\{p\} \times T_p M$  with the tangent space  $T_p M$ . This allows us to see  $T_p M$  as a subset of  $TM$ .

# The Tangent Bundle as a Manifold

## Objective

Let  $M$  be smooth manifold of dimension  $n$ . We would like to bundle together all the tangent spaces  $T_p M$  so as to get a smooth manifold, called the *tangent bundle*.

## Definition

As a set, the *tangent bundle* of  $M$  is the disjoint union,

$$TM := \bigsqcup_{p \in M} T_p M = \{(p, v); p \in M, v \in T_p M\}.$$

## Remarks

- 1 For  $p \in M$  we identify the subset  $\{p\} \times T_p M$  with the tangent space  $T_p M$ . This allows us to see  $T_p M$  as a subset of  $TM$ .
- 2 In particular, we write an element of  $TM$  either as  $(p, v)$  with  $p \in M$  and  $v \in T_p M$ , or simply as  $v$ .

# The Tangent Bundle as a Manifold

## Remark

Let  $U$  be an open set in  $M$ . If  $p \in U$ , then  $T_p U = T_p M$ . Thus,

$$TU = \bigsqcup_{p \in U} T_p U = \bigsqcup_{p \in U} T_p M.$$

# The Tangent Bundle as a Manifold

## Remark

Let  $U$  be an open set in  $M$ . If  $p \in U$ , then  $T_p U = T_p M$ . Thus,

$$TU = \bigsqcup_{p \in U} T_p U = \bigsqcup_{p \in U} T_p M.$$

## Definition

The *canonical map*  $\pi : TM \rightarrow M$  is defined by

$$\pi((p, v)) = p, \quad p \in M, \quad v \in T_p M.$$

# The Tangent Bundle as a Manifold

## Remark

Let  $U$  be an open set in  $M$ . If  $p \in U$ , then  $T_p U = T_p M$ . Thus,

$$TU = \bigsqcup_{p \in U} T_p U = \bigsqcup_{p \in U} T_p M.$$

## Definition

The *canonical map*  $\pi : TM \rightarrow M$  is defined by

$$\pi((p, v)) = p, \quad p \in M, \quad v \in T_p M.$$

## Remarks

- 1 The map  $\pi : TM \rightarrow M$  is onto.



# The Tangent Bundle as a Manifold

## Remark

Let  $U$  be an open set in  $M$ . If  $p \in U$ , then  $T_p U = T_p M$ . Thus,

$$TU = \bigsqcup_{p \in U} T_p U = \bigsqcup_{p \in U} T_p M.$$

## Definition

The *canonical map*  $\pi : TM \rightarrow M$  is defined by

$$\pi((p, v)) = p, \quad p \in M, \quad v \in T_p M.$$

## Remarks

- 1 The map  $\pi : TM \rightarrow M$  is onto.
- 2 If  $p \in M$ , then  $\pi^{-1}(p) = T_p M$ .

# The Tangent Bundle as a Manifold

## Example

Let  $U$  be an open in  $\mathbb{R}^n$ . If  $p \in U$ , then  $T_p U = T_p \mathbb{R}^n = \mathbb{R}^n$ . Recall that, if  $(r^1, \dots, r^n)$  are the standard coordinates on  $\mathbb{R}^n$ , then we identify

$$T_p \mathbb{R}^n \ni v = \sum v^i \frac{\partial}{\partial r^i} \Big|_p \longleftrightarrow \langle v^1, \dots, v^n \rangle \in \mathbb{R}^n.$$

# The Tangent Bundle as a Manifold

## Example

Let  $U$  be an open in  $\mathbb{R}^n$ . If  $p \in U$ , then  $T_p U = T_p \mathbb{R}^n = \mathbb{R}^n$ . Recall that, if  $(r^1, \dots, r^n)$  are the standard coordinates on  $\mathbb{R}^n$ , then we identify

$$T_p \mathbb{R}^n \ni v = \sum v^i \frac{\partial}{\partial r^i} \Big|_p \longleftrightarrow \langle v^1, \dots, v^n \rangle \in \mathbb{R}^n.$$

Thus, the pair  $(p, v)$  is naturally identified with  $(p, v^1, \dots, v^n)$ . Therefore, we have

$$TU = \bigsqcup_{p \in U} T_p U = \bigsqcup_{p \in U} \mathbb{R}^n = U \times \mathbb{R}^n.$$

# The Tangent Bundle as a Manifold

## Aim

We wish to equip  $TM$  with a smooth structure. Therefore, we need to do the following:

# The Tangent Bundle as a Manifold

## Aim

We wish to equip  $TM$  with a smooth structure. Therefore, we need to do the following:

- 1 Define a topology on  $TM$ .

# The Tangent Bundle as a Manifold

## Aim

We wish to equip  $TM$  with a smooth structure. Therefore, we need to do the following:

- 1 Define a topology on  $TM$ .
- 2 Construct a  $C^\infty$ -atlas for  $TM$ .

# The Tangent Bundle as a Manifold

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ .

# The Tangent Bundle as a Manifold

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ .

- Here  $\phi(U)$  is an open in  $\mathbb{R}^n$ , and so  $T(\phi(U)) = \phi(U) \times \mathbb{R}^n$ .



# The Tangent Bundle as a Manifold

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ .

- Here  $\phi(U)$  is an open in  $\mathbb{R}^n$ , and so  $T(\phi(U)) = \phi(U) \times \mathbb{R}^n$ .
- For every  $p \in U$ , the differential  $\phi_{*,p}$  is an isomorphism from  $T_p M = T_p U$  onto  $T_{\phi(p)}(\phi(U)) = \mathbb{R}^n$ .

# The Tangent Bundle as a Manifold

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ .

- Here  $\phi(U)$  is an open in  $\mathbb{R}^n$ , and so  $T(\phi(U)) = \phi(U) \times \mathbb{R}^n$ .
- For every  $p \in U$ , the differential  $\phi_{*,p}$  is an isomorphism from  $T_p M = T_p U$  onto  $T_{\phi(p)}(\phi(U)) = \mathbb{R}^n$ .
- Therefore, we define a map  $\tilde{\phi} : TU \rightarrow \phi(U) \times \mathbb{R}^n$  by

$$\tilde{\phi}(p, v) = (\phi(p), \phi_{*,p}v), \quad p \in U, v \in T_p U.$$

# The Tangent Bundle as a Manifold

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ .

- Here  $\phi(U)$  is an open in  $\mathbb{R}^n$ , and so  $T(\phi(U)) = \phi(U) \times \mathbb{R}^n$ .
- For every  $p \in U$ , the differential  $\phi_{*,p}$  is an isomorphism from  $T_p M = T_p U$  onto  $T_{\phi(p)}(\phi(U)) = \mathbb{R}^n$ .
- Therefore, we define a map  $\tilde{\phi} : TU \rightarrow \phi(U) \times \mathbb{R}^n$  by

$$\tilde{\phi}(p, v) = (\phi(p), \phi_{*,p}v), \quad p \in U, v \in T_p U.$$

- This is a bijection with inverse  $(x, v) \rightarrow (\phi^{-1}(x), \phi_{*,\phi^{-1}(x)}^{-1}v)$ .

# The Tangent Bundle as a Manifold

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ .

- Here  $\phi(U)$  is an open in  $\mathbb{R}^n$ , and so  $T(\phi(U)) = \phi(U) \times \mathbb{R}^n$ .
- For every  $p \in U$ , the differential  $\phi_{*,p}$  is an isomorphism from  $T_p M = T_p U$  onto  $T_{\phi(p)}(\phi(U)) = \mathbb{R}^n$ .

- Therefore, we define a map  $\tilde{\phi} : TU \rightarrow \phi(U) \times \mathbb{R}^n$  by

$$\tilde{\phi}(p, v) = (\phi(p), \phi_{*,p}v), \quad p \in U, v \in T_p U.$$

- This is a bijection with inverse  $(x, v) \rightarrow (\phi^{-1}(x), \phi_{*,\phi^{-1}(x)}^{-1}v)$ .
- This allows us to define a topology on  $TU$  by pulling back the topology of  $\phi(U) \times \mathbb{R}^n$ :

$$W \subseteq TU \text{ is open} \iff \tilde{\phi}(W) \text{ is open in } \phi(U) \times \mathbb{R}^n.$$

# The Tangent Bundle as a Manifold

## Facts

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ .

- Here  $\phi(U)$  is an open in  $\mathbb{R}^n$ , and so  $T(\phi(U)) = \phi(U) \times \mathbb{R}^n$ .
- For every  $p \in U$ , the differential  $\phi_{*,p}$  is an isomorphism from  $T_p M = T_p U$  onto  $T_{\phi(p)}(\phi(U)) = \mathbb{R}^n$ .

- Therefore, we define a map  $\tilde{\phi} : TU \rightarrow \phi(U) \times \mathbb{R}^n$  by

$$\tilde{\phi}(p, v) = (\phi(p), \phi_{*,p}v), \quad p \in U, v \in T_p U.$$

- This is a bijection with inverse  $(x, v) \rightarrow (\phi^{-1}(x), \phi_{*,\phi^{-1}(x)}^{-1}v)$ .
- This allows us to define a topology on  $TU$  by pulling back the topology of  $\phi(U) \times \mathbb{R}^n$ :

$$W \subseteq TU \text{ is open} \iff \tilde{\phi}(W) \text{ is open in } \phi(U) \times \mathbb{R}^n.$$

- With respect to this topology  $\tilde{\phi}$  is a homeomorphism.

# The Tangent Bundle as a Manifold

## Remark

- If  $p \in U$ , then  $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^1} \Big|_p \right\}$  is a basis of  $T_p M$ .

# The Tangent Bundle as a Manifold

## Remark

- If  $p \in U$ , then  $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^1} \Big|_p \right\}$  is a basis of  $T_p M$ .
- The differential  $\phi_{*,p} : T_p U \rightarrow T_{\phi(p)} V = \mathbb{R}^n$  maps  $\frac{\partial}{\partial x^i} \Big|_p$  to  $\frac{\partial}{\partial r^i} \Big|_{\phi(p)}$ . Thus,

$$\sum v^i \frac{\partial}{\partial x^i} \Big|_p \xrightarrow{\phi_*} \sum v^i \frac{\partial}{\partial r^i} \Big|_{\phi(p)} \longleftrightarrow \langle v^1, \dots, v^n \rangle \in \mathbb{R}^n.$$

# The Tangent Bundle as a Manifold

## Remark

- If  $p \in U$ , then  $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$  is a basis of  $T_p M$ .
- The differential  $\phi_{*,p} : T_p U \rightarrow T_{\phi(p)} V = \mathbb{R}^n$  maps  $\frac{\partial}{\partial x^i} \Big|_p$  to  $\frac{\partial}{\partial r^i} \Big|_{\phi(p)}$ . Thus,

$$\sum v^i \frac{\partial}{\partial x^i} \Big|_p \xrightarrow{\phi_*} \sum v^i \frac{\partial}{\partial r^i} \Big|_{\phi(p)} \longleftrightarrow \langle v^1, \dots, v^n \rangle \in \mathbb{R}^n.$$

- If  $\phi(p) = (x^1(p), \dots, x^n(p))$  and  $v = \sum v^i \frac{\partial}{\partial x^i} \Big|_p \in T_p M$ , then  $\tilde{\phi}(p, v) = (\phi(p), \phi_{*,p} v) = (x^1(p), \dots, x^n(p), v^1, \dots, v^n)$ .



# The Tangent Bundle as a Manifold

## Remark

- If  $p \in U$ , then  $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$  is a basis of  $T_p M$ .
- The differential  $\phi_{*,p} : T_p U \rightarrow T_{\phi(p)} V = \mathbb{R}^n$  maps  $\frac{\partial}{\partial x^i} \Big|_p$  to  $\frac{\partial}{\partial r^i} \Big|_{\phi(p)}$ . Thus,

$$\sum v^i \frac{\partial}{\partial x^i} \Big|_p \xrightarrow{\phi_*} \sum v^i \frac{\partial}{\partial r^i} \Big|_{\phi(p)} \longleftrightarrow \langle v^1, \dots, v^n \rangle \in \mathbb{R}^n.$$

- If  $\phi(p) = (x^1(p), \dots, x^n(p))$  and  $v = \sum v^i \frac{\partial}{\partial x^i} \Big|_p \in T_p M$ , then  $\tilde{\phi}(p, v) = (\phi(p), \phi_{*,p} v) = (x^1(p), \dots, x^n(p), v^1, \dots, v^n)$ .

In particular, this defines coordinates on  $TU$ .

# The Tangent Bundle as a Manifold

## Facts

- Let  $(V, \psi) = (V, y^1, \dots, y^n)$  be another chart of  $M$  such that  $U \cap V \neq \emptyset$ . Define  $\tilde{\psi} : TV \rightarrow \psi(V) \times \mathbb{R}^n$  by

$$\tilde{\psi}(p, v) = (\psi(p), \psi_{*,p}v), \quad p \in V, v \in T_pM.$$

# The Tangent Bundle as a Manifold

## Facts

- Let  $(V, \psi) = (V, y^1, \dots, y^n)$  be another chart of  $M$  such that  $U \cap V \neq \emptyset$ . Define  $\tilde{\psi} : TV \rightarrow \psi(V) \times \mathbb{R}^n$  by

$$\tilde{\psi}(p, v) = (\psi(p), \psi_{*,p}v), \quad p \in V, v \in T_pM.$$

- On  $T(U \cap V) = (TU) \cap (TV)$  we have two topologies induced by the topologies of  $TU$  and  $TV$  (defined by the charts  $(U \cap V, \phi|_{U \cap V})$  and  $(U \cap V, \psi|_{U \cap V})$ ).

# The Tangent Bundle as a Manifold

## Facts

- Let  $(V, \psi) = (V, y^1, \dots, y^n)$  be another chart of  $M$  such that  $U \cap V \neq \emptyset$ . Define  $\tilde{\psi} : TV \rightarrow \psi(V) \times \mathbb{R}^n$  by

$$\tilde{\psi}(p, v) = (\psi(p), \psi_{*,p}v), \quad p \in V, v \in T_pM.$$

- On  $T(U \cap V) = (TU) \cap (TV)$  we have two topologies induced by the topologies of  $TU$  and  $TV$  (defined by the charts  $(U \cap V, \phi|_{U \cap V})$  and  $(U \cap V, \psi|_{U \cap V})$ ).

- On  $\tilde{\phi}(TU \cap TV) = \phi(U \cap V) \times \mathbb{R}^n$  we have

$$\tilde{\psi} \circ \tilde{\phi}^{-1}(r, v) = (\psi \circ \phi^{-1}(r), \psi_* \circ \phi_*^{-1}v) = (\psi \circ \phi^{-1}(r), (\psi \circ \phi^{-1})_{*,r}v).$$

# The Tangent Bundle as a Manifold

## Facts

- Let  $(V, \psi) = (V, y^1, \dots, y^n)$  be another chart of  $M$  such that  $U \cap V \neq \emptyset$ . Define  $\tilde{\psi} : TV \rightarrow \psi(V) \times \mathbb{R}^n$  by

$$\tilde{\psi}(p, v) = (\psi(p), \psi_{*,p}v), \quad p \in V, v \in T_pM.$$

- On  $T(U \cap V) = (TU) \cap (TV)$  we have two topologies induced by the topologies of  $TU$  and  $TV$  (defined by the charts  $(U \cap V, \phi|_{U \cap V})$  and  $(U \cap V, \psi|_{U \cap V})$ ).

- On  $\tilde{\phi}(TU \cap TV) = \phi(U \cap V) \times \mathbb{R}^n$  we have

$$\tilde{\psi} \circ \tilde{\phi}^{-1}(r, v) = (\psi \circ \phi^{-1}(r), \psi_* \circ \phi_*^{-1}v) = (\psi \circ \phi^{-1}(r), (\psi \circ \phi^{-1})_{*,r}v).$$

- Here  $(\psi \circ \phi^{-1})_{*,r}$  is the multiplication by the Jacobian matrix  $J_{\psi \circ \phi^{-1}}(r) = [\partial(y^j \circ \phi^{-1})/\partial r^i(r)]$  whose entries depends smoothly on  $r$ .

# The Tangent Bundle as a Manifold

## Facts

- Let  $(V, \psi) = (V, y^1, \dots, y^n)$  be another chart of  $M$  such that  $U \cap V \neq \emptyset$ . Define  $\tilde{\psi} : TV \rightarrow \psi(V) \times \mathbb{R}^n$  by

$$\tilde{\psi}(p, v) = (\psi(p), \psi_{*,p}v), \quad p \in V, v \in T_pM.$$

- On  $T(U \cap V) = (TU) \cap (TV)$  we have two topologies induced by the topologies of  $TU$  and  $TV$  (defined by the charts  $(U \cap V, \phi|_{U \cap V})$  and  $(U \cap V, \psi|_{U \cap V})$ ).

- On  $\tilde{\phi}(TU \cap TV) = \phi(U \cap V) \times \mathbb{R}^n$  we have

$$\tilde{\psi} \circ \tilde{\phi}^{-1}(r, v) = (\psi \circ \phi^{-1}(r), \psi_* \circ \phi_*^{-1}v) = (\psi \circ \phi^{-1}(r), (\psi \circ \phi^{-1})_{*,r}v).$$

- Here  $(\psi \circ \phi^{-1})_{*,r}$  is the multiplication by the Jacobian matrix  $J_{\psi \circ \phi^{-1}}(r) = [\partial(y^j \circ \phi^{-1})/\partial r^i(r)]$  whose entries depends smoothly on  $r$ .

- Therefore,  $\tilde{\psi} \circ \tilde{\phi}^{-1} : \phi(U \cap V) \times \mathbb{R}^n \rightarrow \psi(U \cap V) \times \mathbb{R}^n$  is smooth map. Its inverse map  $\tilde{\phi} \circ \tilde{\psi}^{-1}$  is smooth as well, and so  $\tilde{\psi} \circ \tilde{\phi}^{-1}$  is a diffeomorphism.

# The Tangent Bundle as a Manifold

## Facts (Continued)

- $T(U \cap V)$  is open in  $TU$  and in  $TV$ .

# The Tangent Bundle as a Manifold

## Facts (Continued)

- $T(U \cap V)$  is open in  $TU$  and in  $TV$ .
- As  $\tilde{\psi} \circ \tilde{\phi}^{-1} : \phi(U \cap V) \times \mathbb{R}^n \rightarrow \psi(U \cap V) \times \mathbb{R}^n$  is a diffeomorphism, this is a homeomorphism.



# The Tangent Bundle as a Manifold

## Facts (Continued)

- $T(U \cap V)$  is open in  $TU$  and in  $TV$ .
- As  $\tilde{\psi} \circ \tilde{\phi}^{-1} : \phi(U \cap V) \times \mathbb{R}^n \rightarrow \psi(U \cap V) \times \mathbb{R}^n$  is a diffeomorphism, this is a homeomorphism.
- If  $W \subseteq T(U \cap V)$ , then
$$\begin{aligned} W \text{ open in } TU &\iff \tilde{\phi}(W) \text{ open in } \phi(U) \times \mathbb{R}^n, \\ &\iff \tilde{\psi} \circ \tilde{\phi}^{-1}[\tilde{\phi}(W)] \text{ open in } \psi(U \cap V) \times \mathbb{R}^n, \\ &\iff \tilde{\psi}(W) \text{ open in } \psi(V) \times \mathbb{R}^n, \\ &\iff W \text{ open in } TV. \end{aligned}$$

Thus,  $TU$  and  $TV$  induce the same topology on  $T(U \cap V)$ .

# The Tangent Bundle as a Manifold

## Facts (Continued)

- $T(U \cap V)$  is open in  $TU$  and in  $TV$ .
- As  $\tilde{\psi} \circ \tilde{\phi}^{-1} : \phi(U \cap V) \times \mathbb{R}^n \rightarrow \psi(U \cap V) \times \mathbb{R}^n$  is a diffeomorphism, this is a homeomorphism.
- If  $W \subseteq T(U \cap V)$ , then
$$\begin{aligned} W \text{ open in } TU &\iff \tilde{\phi}(W) \text{ open in } \phi(U) \times \mathbb{R}^n, \\ &\iff \tilde{\psi} \circ \tilde{\phi}^{-1}[\tilde{\phi}(W)] \text{ open in } \psi(U \cap V) \times \mathbb{R}^n, \\ &\iff \tilde{\psi}(W) \text{ open in } \psi(V) \times \mathbb{R}^n, \\ &\iff W \text{ open in } TV. \end{aligned}$$

Thus,  $TU$  and  $TV$  induce the same topology on  $T(U \cap V)$ .

- It follows that, if  $W$  is open in  $TU$  and  $X$  is open in  $TV$ , then  $W \cap X = (W \cap T(U \cap V)) \cap (X \cap T(U \cap V))$  is open in  $T(U \cap V)$  (by definition of the subspace topology).

# The Tangent Bundle as a Manifold

## Summary

- (a) If  $(U, \phi)$  is a chart for  $M$ , then  $\tilde{\phi} : TU \rightarrow \phi(U) \times \mathbb{R}^n$  allows us to define a topology on  $TU$  by pulling back the topology of  $\phi(U) \times \mathbb{R}^n$ . This map then becomes a homeomorphism.

# The Tangent Bundle as a Manifold

## Summary

- (a) If  $(U, \phi)$  is a chart for  $M$ , then  $\tilde{\phi} : TU \rightarrow \phi(U) \times \mathbb{R}^n$  allows us to define a topology on  $TU$  by pulling back the topology of  $\phi(U) \times \mathbb{R}^n$ . This map then becomes a homeomorphism.
- (b) If  $(V, \psi)$  is another chart for  $M$ , then the transition map  $\tilde{\psi} \circ \tilde{\phi}^{-1} : \phi(U \cap V) \times \mathbb{R}^n \rightarrow \psi(U \cap V) \times \mathbb{R}^n$  is a diffeomorphism.

# The Tangent Bundle as a Manifold

## Summary

- (a) If  $(U, \phi)$  is a chart for  $M$ , then  $\tilde{\phi} : TU \rightarrow \phi(U) \times \mathbb{R}^n$  allows us to define a topology on  $TU$  by pulling back the topology of  $\phi(U) \times \mathbb{R}^n$ . This map then becomes a homeomorphism.
- (b) If  $(V, \psi)$  is another chart for  $M$ , then the transition map  $\tilde{\psi} \circ \tilde{\phi}^{-1} : \phi(U \cap V) \times \mathbb{R}^n \rightarrow \psi(U \cap V) \times \mathbb{R}^n$  is a diffeomorphism.
- (c)  $TU$  and  $TV$  induce the same topology on  $T(U \cap V)$ . In particular, if  $W$  is open in  $TU$  and  $X$  is open in  $TV$ , then  $W \cap X = (W \cap T(U \cap V)) \cap (X \cap T(U \cap V))$  is open in  $T(U \cap V)$ .

In particular, (b) would allow us to get a  $C^\infty$  atlas for  $TM$  provided we can define a topology on  $TM$  by patching together the  $TU$ -topologies.

# The Tangent Bundle as a Manifold

## Reminder (Topological Bases; see Appendix A)

Let  $X$  be a topological space. A *basis for the topology of  $X$*  is a collection  $\mathcal{B}$  of open sets such that, for every open  $U \subseteq X$  and every  $p \in U$ , there is an open set  $V \in \mathcal{B}$  such that  $p \in V \subseteq U$ .

# The Tangent Bundle as a Manifold

## Reminder (Topological Bases; see Appendix A)

Let  $X$  be a topological space. A *basis for the topology of  $X$*  is a collection  $\mathcal{B}$  of open sets such that, for every open  $U \subseteq X$  and every  $p \in U$ , there is an open set  $V \in \mathcal{B}$  such that  $p \in V \subseteq U$ .

## Remark

If  $\mathcal{B}$  is a basis for the topology of  $X$ , then every open set is the union of open sets in  $\mathcal{B}$ . We then say that  $\mathcal{B}$  generates the topology of  $X$ .

# The Tangent Bundle as a Manifold

## Proposition (Proposition A.8)

Let  $X$  be a set and  $\mathcal{B}$  a collection of subsets such that:

(i)  $X = \cup_{V \in \mathcal{B}} V$ .



# The Tangent Bundle as a Manifold

## Proposition (Proposition A.8)

Let  $X$  be a set and  $\mathcal{B}$  a collection of subsets such that:

- (i)  $X = \bigcup_{V \in \mathcal{B}} V$ .
- (ii) If  $V_1, V_2 \in \mathcal{B}$  and  $p \in V_1 \cap V_2$ , then there is  $W \in \mathcal{B}$  such that  $p \in W \subseteq V_1 \cap V_2$ .

Then:

# The Tangent Bundle as a Manifold

## Proposition (Proposition A.8)

Let  $X$  be a set and  $\mathcal{B}$  a collection of subsets such that:

- (i)  $X = \bigcup_{V \in \mathcal{B}} V$ .
- (ii) If  $V_1, V_2 \in \mathcal{B}$  and  $p \in V_1 \cap V_2$ , then there is  $W \in \mathcal{B}$  such that  $p \in W \subseteq V_1 \cap V_2$ .

Then:

- 1  $\mathcal{B}$  is a basis for a unique topology on  $X$ .

# The Tangent Bundle as a Manifold

## Proposition (Proposition A.8)

Let  $X$  be a set and  $\mathcal{B}$  a collection of subsets such that:

- (i)  $X = \bigcup_{V \in \mathcal{B}} V$ .
- (ii) If  $V_1, V_2 \in \mathcal{B}$  and  $p \in V_1 \cap V_2$ , then there is  $W \in \mathcal{B}$  such that  $p \in W \subseteq V_1 \cap V_2$ .

Then:

- ①  $\mathcal{B}$  is a basis for a unique topology on  $X$ .
- ② The open sets for this topology consists of unions of sets in  $\mathcal{B}$ .

# The Tangent Bundle as a Manifold

## Proposition (Proposition A.8)

Let  $X$  be a set and  $\mathcal{B}$  a collection of subsets such that:

- (i)  $X = \bigcup_{V \in \mathcal{B}} V$ .
- (ii) If  $V_1, V_2 \in \mathcal{B}$  and  $p \in V_1 \cap V_2$ , then there is  $W \in \mathcal{B}$  such that  $p \in W \subseteq V_1 \cap V_2$ .

Then:

- ①  $\mathcal{B}$  is a basis for a unique topology on  $X$ .
- ② The open sets for this topology consists of unions of sets in  $\mathcal{B}$ .

## Remark

The condition (ii) holds automatically if  $\mathcal{B}$  is closed under finite intersection.

# The Tangent Bundle as a Manifold

## Facts

Let  $\mathcal{A} = \{(U_\alpha, \phi_\alpha)\}$  be the maximal atlas of  $M$  (which defines its smooth structure). Define

$$\mathcal{B} = \bigcup_{\alpha} \{W; W \text{ is an open in } TU_\alpha\}.$$

Note that  $TU_\alpha \in \mathcal{B}$ .

# The Tangent Bundle as a Manifold

## Facts

Let  $\mathcal{A} = \{(U_\alpha, \phi_\alpha)\}$  be the maximal atlas of  $M$  (which defines its smooth structure). Define

$$\mathcal{B} = \bigcup_{\alpha} \{W; W \text{ is an open in } TU_\alpha\}.$$

Note that  $TU_\alpha \in \mathcal{B}$ .

- As  $\bigcup U_\alpha = M$ , we have

$$\bigcup_{\alpha} TU_\alpha = \bigcup_{\alpha} \bigsqcup_{p \in U_\alpha} T_p M = \bigsqcup_{p \in \bigcup U_\alpha} T_p M = \bigsqcup_{p \in M} T_p M = TM.$$

# The Tangent Bundle as a Manifold

## Facts

Let  $\mathcal{A} = \{(U_\alpha, \phi_\alpha)\}$  be the maximal atlas of  $M$  (which defines its smooth structure). Define

$$\mathcal{B} = \bigcup_{\alpha} \{W; W \text{ is an open in } TU_\alpha\}.$$

Note that  $TU_\alpha \in \mathcal{B}$ .

- As  $\bigcup_{\alpha} U_\alpha = M$ , we have

$$\bigcup_{\alpha} TU_\alpha = \bigcup_{\alpha} \bigsqcup_{p \in U_\alpha} T_p M = \bigsqcup_{p \in \bigcup_{\alpha} U_\alpha} T_p M = \bigsqcup_{p \in M} T_p M = TM.$$

- If  $W_\alpha$  is an open in  $TU_\alpha$  and  $W_\beta$  is an open in  $TU_\beta$ , then  $W_1 \cap W_2$  is open in  $T(U_\alpha \cap U_\beta)$ , and hence is contained in  $\mathcal{B}$ .

# The Tangent Bundle as a Manifold

## Facts

Let  $\mathcal{A} = \{(U_\alpha, \phi_\alpha)\}$  be the maximal atlas of  $M$  (which defines its smooth structure). Define

$$\mathcal{B} = \bigcup_{\alpha} \{W; W \text{ is an open in } TU_\alpha\}.$$

Note that  $TU_\alpha \in \mathcal{B}$ .

- As  $\bigcup U_\alpha = M$ , we have

$$\bigcup_{\alpha} TU_\alpha = \bigcup_{\alpha} \bigsqcup_{p \in U_\alpha} T_p M = \bigsqcup_{p \in \bigcup U_\alpha} T_p M = \bigsqcup_{p \in M} T_p M = TM.$$

- If  $W_\alpha$  is an open in  $TU_\alpha$  and  $W_\beta$  is an open in  $TU_\beta$ , then  $W_1 \cap W_2$  is open in  $T(U_\alpha \cap U_\beta)$ , and hence is contained in  $\mathcal{B}$ .

It follows that  $\mathcal{B}$  satisfies the conditions (i) and (ii) of Proposition A.8, and so it's a basis for a unique topology on  $TM$ .



# The Tangent Bundle as a Manifold

## Definition

The topology of  $TM$  is the topology generated by  $\mathcal{B}$ . The open sets are unions of sets in  $\mathcal{B}$ .

# The Tangent Bundle as a Manifold

## Definition

The topology of  $TM$  is the topology generated by  $\mathcal{B}$ . The open sets are unions of sets in  $\mathcal{B}$ .

## Remark

Each  $TU_\alpha$  is open in  $TM$ , since it is contained in  $\mathcal{B}$ .

# The Tangent Bundle as a Manifold

## Definition

The topology of  $TM$  is the topology generated by  $\mathcal{B}$ . The open sets are unions of sets in  $\mathcal{B}$ .

## Remark

Each  $TU_\alpha$  is open in  $TM$ , since it is contained in  $\mathcal{B}$ .

## Proposition (Proposition 12.4)

*As a topological space  $TM$  is Hausdorff.*

# The Tangent Bundle as a Manifold

## Definition

The topology of  $TM$  is the topology generated by  $\mathcal{B}$ . The open sets are unions of sets in  $\mathcal{B}$ .

## Remark

Each  $TU_\alpha$  is open in  $TM$ , since it is contained in  $\mathcal{B}$ .

## Proposition (Proposition 12.4)

*As a topological space  $TM$  is Hausdorff.*

## Remark

Each open  $TU_\alpha$  is Hausdorff since it is homeomorphic to the open set  $\phi_\alpha(U) \times \mathbb{R}^n \subseteq \mathbb{R}^n \times \mathbb{R}^n$ . This can be used to show that  $TM$  is Hausdorff.

# The Tangent Bundle as a Manifold

## Proposition (Proposition 12.3)

*The topology of  $TM$  is second countable.*

# The Tangent Bundle as a Manifold

## Proposition (Proposition 12.3)

*The topology of  $TM$  is second countable.*

## Remarks

- It can be shown that the topology of  $M$  admits a countable basis  $\{U_i\}_{i \in I}$  consisting of domains of charts (cf. Lemma 12.2 of Tu's book).

# The Tangent Bundle as a Manifold

## Proposition (Proposition 12.3)

*The topology of  $TM$  is second countable.*

## Remarks

- It can be shown that the topology of  $M$  admits a countable basis  $\{U_i\}_{i \in I}$  consisting of domains of charts (cf. Lemma 12.2 of Tu's book).
- Each  $TU_i$  is second countable since it is homeomorphic to an open in  $\mathbb{R}^n \times \mathbb{R}^n$ .

# The Tangent Bundle as a Manifold

## Proposition (Proposition 12.3)

*The topology of  $TM$  is second countable.*

## Remarks

- It can be shown that the topology of  $M$  admits a countable basis  $\{U_i\}_{i \in I}$  consisting of domains of charts (cf. Lemma 12.2 of Tu's book).
- Each  $TU_i$  is second countable since it is homeomorphic to an open in  $\mathbb{R}^n \times \mathbb{R}^n$ .
- If for each  $i \in I$ , we let  $\{W_{i,j}\}_{j \in \mathbb{N}}$  be a countable basis for the topology of  $TU_i$ , then  $\{W_{i,j}; i \in I, j \in \mathbb{N}\}$  is a countable basis for the topology of  $TM$  (see Tu's book).



# The Tangent Bundle as a Manifold

## Facts

- Each  $TU_\alpha$  is an open in  $TM$ .

# The Tangent Bundle as a Manifold

## Facts

- Each  $TU_\alpha$  is an open in  $TM$ .
- We know that  $TM = \bigcup_\alpha TU_\alpha$ .

# The Tangent Bundle as a Manifold

## Facts

- Each  $TU_\alpha$  is an open in  $TM$ .
- We know that  $TM = \bigcup_\alpha TU_\alpha$ .
- The local trivializations  $\tilde{\phi}_\alpha : TU_\alpha \rightarrow \phi_\alpha(U_\alpha) \times \mathbb{R}^n$  are homeomorphisms onto open sets in  $\mathbb{R}^n \times \mathbb{R}^n$ .

# The Tangent Bundle as a Manifold

## Facts

- Each  $TU_\alpha$  is an open in  $TM$ .
- We know that  $TM = \bigcup_\alpha TU_\alpha$ .
- The local trivializations  $\tilde{\phi}_\alpha : TU_\alpha \rightarrow \phi_\alpha(U_\alpha) \times \mathbb{R}^n$  are homeomorphisms onto open sets in  $\mathbb{R}^n \times \mathbb{R}^n$ .
- All the transition maps  $\tilde{\phi}_\beta \circ \tilde{\phi}_\alpha^{-1}$  are smooth.

# The Tangent Bundle as a Manifold

## Facts

- Each  $TU_\alpha$  is an open in  $TM$ .
- We know that  $TM = \bigcup_\alpha TU_\alpha$ .
- The local trivializations  $\tilde{\phi}_\alpha : TU_\alpha \rightarrow \phi_\alpha(U_\alpha) \times \mathbb{R}^n$  are homeomorphisms onto open sets in  $\mathbb{R}^n \times \mathbb{R}^n$ .
- All the transition maps  $\tilde{\phi}_\beta \circ \tilde{\phi}_\alpha^{-1}$  are smooth.

## Proposition

*The collection  $\{(TU_\alpha, \tilde{\phi}_\alpha)\}$  is a  $C^\infty$  atlas for  $TM$ , and hence  $TM$  is a smooth manifold of dimension  $2n$ .*

# The Tangent Bundle as a Manifold

## Facts

- Each  $TU_\alpha$  is an open in  $TM$ .
- We know that  $TM = \bigcup_\alpha TU_\alpha$ .
- The local trivializations  $\tilde{\phi}_\alpha : TU_\alpha \rightarrow \phi_\alpha(U_\alpha) \times \mathbb{R}^n$  are homeomorphisms onto open sets in  $\mathbb{R}^n \times \mathbb{R}^n$ .
- All the transition maps  $\tilde{\phi}_\beta \circ \tilde{\phi}_\alpha^{-1}$  are smooth.

## Proposition

*The collection  $\{(TU_\alpha, \tilde{\phi}_\alpha)\}$  is a  $C^\infty$  atlas for  $TM$ , and hence  $TM$  is a smooth manifold of dimension  $2n$ .*

## Remark

If  $\{(V_\beta, \psi_\beta)\}$  is any  $C^\infty$ -atlas for  $M$ , then we also get a  $C^\infty$  atlas  $\{(TV_\beta, \tilde{\psi}_\beta)\}$  for  $TM$ . It is compatible with the atlas  $\{(TU_\alpha, \tilde{\phi}_\alpha)\}$ , and so it defines the same smooth structure.

# The Tangent Bundle as a Manifold

## Facts

- The canonical map  $\pi : TM \rightarrow M$  is such that  $\pi(v) = p$  if  $v \in T_pM$ . It is onto.

# The Tangent Bundle as a Manifold

## Facts

- The canonical map  $\pi : TM \rightarrow M$  is such that  $\pi(v) = p$  if  $v \in T_p M$ . It is onto.
- Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ . Then

$$\begin{aligned}\phi \circ \pi \circ \tilde{\phi}^{-1}(r^1, \dots, r^n, v^1, \dots, v^n) \\&= \phi \circ \pi \left[ \phi^{-1}(r^1, \dots, r^n), \sum v^i \partial / \partial x^i \right] \\&= \phi \circ \phi^{-1}(r^1, \dots, r^n) = (r^1, \dots, r^n).\end{aligned}$$



# The Tangent Bundle as a Manifold

## Facts

- The canonical map  $\pi : TM \rightarrow M$  is such that  $\pi(v) = p$  if  $v \in T_p M$ . It is onto.
- Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ . Then

$$\begin{aligned}\phi \circ \pi \circ \tilde{\phi}^{-1}(r^1, \dots, r^n, v^1, \dots, v^n) \\&= \phi \circ \pi \left[ \phi^{-1}(r^1, \dots, r^n), \sum v^i \partial / \partial x^i \right] \\&= \phi \circ \phi^{-1}(r^1, \dots, r^n) = (r^1, \dots, r^n).\end{aligned}$$

- As  $(U, \phi)$  and  $(TU, \tilde{\phi})$  are charts this shows that  $\pi$  is  $C^\infty$ .

# The Tangent Bundle as a Manifold

## Facts

- The canonical map  $\pi : TM \rightarrow M$  is such that  $\pi(v) = p$  if  $v \in T_p M$ . It is onto.
- Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ . Then

$$\begin{aligned}\phi \circ \pi \circ \tilde{\phi}^{-1}(r^1, \dots, r^n, v^1, \dots, v^n) \\&= \phi \circ \pi \left[ \phi^{-1}(r^1, \dots, r^n), \sum v^i \partial / \partial x^i \right] \\&= \phi \circ \phi^{-1}(r^1, \dots, r^n) = (r^1, \dots, r^n).\end{aligned}$$

- As  $(U, \phi)$  and  $(TU, \tilde{\phi})$  are charts this shows that  $\pi$  is  $C^\infty$ .
- By the converse of the submersion theorem (exercise!) this also shows that  $\pi$  is a submersion.

# The Tangent Bundle as a Manifold

## Facts

- The canonical map  $\pi : TM \rightarrow M$  is such that  $\pi(v) = p$  if  $v \in T_p M$ . It is onto.
- Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart for  $M$ . Then

$$\begin{aligned}\phi \circ \pi \circ \tilde{\phi}^{-1}(r^1, \dots, r^n, v^1, \dots, v^n) \\&= \phi \circ \pi \left[ \phi^{-1}(r^1, \dots, r^n), \sum v^i \partial / \partial x^i \right] \\&= \phi \circ \phi^{-1}(r^1, \dots, r^n) = (r^1, \dots, r^n).\end{aligned}$$

- As  $(U, \phi)$  and  $(TU, \tilde{\phi})$  are charts this shows that  $\pi$  is  $C^\infty$ .
- By the converse of the submersion theorem (exercise!) this also shows that  $\pi$  is a submersion.

## Proposition

*The canonical projection  $\pi : TM \rightarrow M$  is a surjective submersion.*