

Noncommutative Geometry  
Chapter 1:  
Spectrum and Duality Spaces/Algebras

Sichuan University, Spring 2025

## Main References

- Connes, A.: *Noncommutative geometry*. Academic Press, San Diego, 1994.
- Gracia-Bondia, J.M.; Varilly, J.C.; Figueroa, H.: *Elements of Noncommutative Geometry*. Birkhäuser, Boston, 2001.
- Lectures notes to be posted online.

## References for this Chapter

- Arveson, W.: *A Short Course on Spectral Theory*. Graduate Texts in Mathematics, Springer, 2002 .
- Gracia-Bondia, J.M.; Varilly, J.C.; Figueroa, H.: *Elements of Noncommutative Geometry*.

## Remarks

- All the vector spaces and algebras are vector spaces or algebras over  $\mathbb{C}$ .
- Unless otherwise mentioned all the topological spaces are Hausdorff.

## Definition

A Banach algebra is an algebra  $A$  endowed with a Banach norm  $\|\cdot\|$  such that

$$\|xy\| \leq \|x\|\|y\| \quad \forall x, y \in A.$$

## Definition

A  $C^*$ -algebra is a Banach algebra  $A$  together with an antilinear involution  $x \rightarrow x^*$  such that

$$\begin{aligned} (xy)^* &= y^*x^* & \forall x, y \in A, \\ \|x^*\| &= \|x\| \quad \text{and} \quad \|x^*x\| = \|x\|^2 & \forall x \in A. \end{aligned}$$

# \*-Homomorphisms

## Definition

Let  $A$  and  $B$  be  $C^*$ -algebras.

- 1 A  $*$ -homomorphism  $\phi : A \rightarrow B$  is a continuous homomorphism of algebras such that  $\phi(x^*) = \phi(x)^*$  for all  $x \in A$ .
- 2 A  $*$ -isomorphism  $\phi : A \rightarrow B$  is  $*$ -homomorphism which is bijective.

## Remarks

- By the open mapping theorem any bijective continuous linear map between Banach spaces has a continuous inverse. Therefore, the inverse of any  $*$ -isomorphism is continuous.
- It can be shown that any  $*$ -isomorphism between  $C^*$ -algebras is isometric.

## Example

Let  $M_n(\mathbb{C})$  be the algebra of  $n \times n$ -matrices with complex entries.

- Its involution is  $A \rightarrow A^*$ , where  $A^*$  is the adjoint of  $A$ .
- A  $C^*$ -algebra norm is given by the norm defined by

$$\|A\| := \sup\{\|Ax\|; x \in \mathbb{C}^n, \|x\| = 1\}, \quad A \in M_n(\mathbb{C}).$$

## Example

Let  $\mathcal{H}$  be a Hilbert space and  $\mathcal{L}(\mathcal{H})$  the algebra of continuous linear operators  $T : \mathcal{H} \rightarrow \mathcal{H}$ .

- The involution of  $\mathcal{L}(\mathcal{H})$  is  $T \rightarrow T^*$ , where  $T^*$  is the adjoint of  $T$ , i.e., the unique linear operator on  $\mathcal{H}$  such that

$$\langle T^* \xi, \eta \rangle = \langle \xi, T \eta \rangle \quad \forall \xi, \eta \in \mathcal{H}.$$

- The norm of  $\mathcal{L}(\mathcal{H})$  is

$$\|T\| := \sup_{\|\xi\|=1} \|T\xi\|$$

- More generally, any closed  $*$ -subalgebra of  $\mathcal{L}(\mathcal{H})$  is a  $C^*$ -algebra.

# \*-Representations

## Definition

A  $*$ -representation of a  $C^*$ -algebra in a Hilbert space  $\mathcal{H}$  is a  $*$ -homomorphism from  $A$  to  $\mathcal{L}(\mathcal{H})$ .

## Theorem (Gel'fand-Naimark)

*Any  $C^*$ -algebra  $A$  admits an isometric  $*$ -representation  $\pi$  in some Hilbert space  $\mathcal{H}$ .*

## Remark

Any isometric linear map between Banach spaces has closed range and is an isomorphism onto its range (Exercise!).

## Consequence

Any  $C^*$ -algebra  $A$  can be  $*$ -represented as a closed  $*$ -subalgebra of some  $\mathcal{L}(\mathcal{H})$ .



## Example

Let  $X$  be a compact (Hausdorff) space and  $C(X)$  its algebra of continuous complex-valued functions.

- The involution of  $C(X)$  is  $f \rightarrow \bar{f}$ , where  $\bar{f}$  is the complex conjugate of  $f$ .
- The norm of  $C(X)$  is

$$\|f\|_{C(X)} := \sup_{x \in X} |f(x)|, \quad f \in C(X).$$

- The constant function  $1$  is a unit for  $C(X)$ , and so  $C(X)$  is a unital commutative  $C^*$ -algebra.

# Further Examples

## Example

Let  $X$  be a locally compact topological space and  $C_0(X)$  its algebra of continuous functions “vanishing at infinity”.

- Recall that  $f \in C(X)$  vanishes at infinity iff, for all  $\epsilon > 0$ , there  $K \subset X$  compact s.t.  $|f(x)| \leq \epsilon$  on  $X \setminus K$ .
- The involution of  $C_0(X)$  is  $f \rightarrow \bar{f}$ .
- The norm of  $C_0(X)$  is

$$\|f\|_{C_0(X)} := \sup_{x \in X} |f(x)|, \quad f \in C_0(X).$$

- The  $C^*$ -algebra  $C_0(X)$  is commutative, but it is not unital, since  $1 \notin C_0(X)$ .

## Remark

It can be shown that  $C(X)$  and  $C_0(X)$  are essentially the only examples of commutative  $C^*$ -algebras.

# Adding a Unit

## Definition

Let  $A$  be a (possibly non-unital) Banach algebra. Define

$$A^+ = A \oplus \mathbb{C}.$$

We endow  $A^+$  with the product and norm given by

$$(x_1, \lambda_1) \cdot (x_2, \lambda_2) := (x_1 x_2 + \lambda_1 x_2 + \lambda_2 x_1, \lambda_1 \lambda_2), \quad (x_j, \lambda_j) \in A^+, \\ \| (x, \lambda) \|_{A^+} := \sup \{ \| xy + \lambda y \|_A; y \in A, \| y \|_A = 1 \}, \quad (x, \lambda) \in A^+.$$

## Proposition

- 1  $A^+$  is a Banach algebra with unit  $1_{A^+} := (0, 1)$ .
- 2 The map  $x \rightarrow (x, 0)$  is an isometric embedding of  $A$  into  $A^+$ .

## Definition

$A^+$  is called the unitalization of  $A$ .

# Adding a Unit

## Remark

- The embedding of  $x \rightarrow (x, 0)$  identifies  $A$  with the closed ideal  $A \oplus \{0\}$  of  $A^+$ .
- This allows us to write any element of  $A^+$  as

$$(x, \lambda) = x + \lambda 1_{A^+}, \quad x \in A, \quad \lambda \in \mathbb{C}.$$

## Proposition

Assume  $A$  is a  $C^*$ -algebra, and equip  $A^+$  with the involution

$$(x + \lambda 1_{A^+})^* = x^* + \bar{\lambda} 1_{A^+}, \quad x \in A, \quad \lambda \in \mathbb{C}.$$

Then  $A$  is a  $C^*$ -algebra.

## Example

Let  $A = C_0(\Omega)$ , where  $\Omega \subset \mathbb{R}^n$  is a bounded open set with boundary  $\partial\Omega$ .

- Here

$$C_0(\Omega) = \{f \in C(\overline{\Omega}); f = 0 \text{ on } \partial\Omega\}.$$

- The unitalization of  $A$  is

$$A^+ = \{f \in C(\overline{\Omega}); f|_{\partial\Omega} \text{ is constant}\}.$$

## Setup

- $A$  is a Banach algebra with a unit  $1_A$  such that  $\|1_A\| = 1$ .
- This is always satisfied if  $A$  is a  $C^*$ -algebra.
- $A^{-1}$  is the group of invertible elements of  $A$ . This is an open subset of  $A$ .

## Definition

Let  $x \in A$ . The spectrum of  $x$  is

$$\operatorname{Sp}_A(x) := \{\lambda \in \mathbb{C}; x - \lambda \notin A^{-1}\}.$$

The complement of  $\operatorname{Sp}_A(x)$  is called the resolvent set of  $x$ .

## Proposition

- 1  $\text{Sp}_A(x)$  is a non-empty compact subset of  $\mathbb{C}$  contained in the disk  $\overline{D(0, \|x\|)}$ .
- 2 The resolvent  $\lambda \rightarrow (x - \lambda)^{-1}$  is an analytic map from  $\mathbb{C} \setminus \text{Sp}_A(x)$  to  $A$ .

## Partial Proof.

- By definition  $\lambda \in \mathbb{C} \setminus \text{Sp}_A(x) \Leftrightarrow x - \lambda \in A^{-1}$ .
- Here  $\mathbb{C} \ni \lambda \rightarrow x - \lambda \in A$  is continuous and  $A^{-1}$  is an open set of  $A$ .
- Thus,  $\mathbb{C} \setminus \text{Sp}_A(x)$  is an open set, and hence  $\text{Sp}_A(x)$  is closed.



## Partial Proof (Continued).

- If  $\|x\| < 1$ , then  $\sum_{n \geq 0} x^n = (1 - x)^{-1}$ .
- If  $\lambda > \|x\|$ , then  $\lambda^{-1}\|x\| < 1$ , and so

$$\sum \lambda^{-n-1} x^n = \lambda^{-1} \sum (\lambda^{-1} x)^n = \lambda^{-1} (1 - \lambda^{-1} x)^{-1} = (\lambda - x)^{-1}.$$

In particular,  $x - \lambda \in A^{-1}$ , i.e.,  $\lambda \in \mathbb{C} \setminus \text{Sp}_A(x)$ .

- Thus,  $\mathbb{C} \setminus \overline{D}(0, \|x\|) \subset \mathbb{C} \setminus \text{Sp}_A(x)$ , i.e.,  $\text{Sp}_A(x) \subset \overline{D}(0, \|x\|)$ .
- Therefore,  $\text{Sp}_A(x)$  is a bounded closed set, and so this is a compact set.





## Remarks

- If  $A$  is not unital, we define the spectrum of  $x \in A$  to be its spectrum in  $A^+$ , i.e.,

$$\mathrm{Sp}_A(x) := \mathrm{Sp}_{A^+}(x).$$

- $0$  is always contained in  $\mathrm{Sp}_{A^+}(x)$ , since the proper ideal  $A$  cannot contain invertible elements of  $A^+$ .
- If  $A$  is unital, then  $\mathrm{Sp}_{A^+}(x) = \mathrm{Sp}_A(x) \cup \{0\}$ .

## Example

Let  $A = C(X)$ , where  $X$  is a compact (Hausdorff) space.

- If  $f \in C(X)$  and  $\lambda \in \mathbb{C}$ , then

$$f - \lambda \text{ invertible} \iff (f(x) - \lambda \neq 0 \ \forall x \in X) \iff \lambda \notin f(X).$$

- Thus,  $\mathbb{C} \setminus \text{Sp}_{C(X)}(f) = \mathbb{C} \setminus f(X)$ , i.e.,

$$\text{Sp}_{C(X)}(f) = f(X).$$

## Example

Let  $A = C_0(\Omega)$ , where  $\Omega \subset \mathbb{R}^n$  is a bounded open set.

- Here  $A^+ = \{f \in C(\overline{\Omega}); f|_{\partial\Omega} \text{ is constant} \}$ .
- Thus, if  $f \in C_0(\Omega)$ , then, by spectral permanence,

$$\text{Sp}_{C_0(\Omega)}(f) = \text{Sp}_{A^+}(f) = \text{Sp}_{C(\overline{\Omega})}(f) = f(\overline{\Omega}) = f(\Omega) \cup \{0\}.$$

- More generally, if  $X$  is a locally compact Hausdorff space, then

$$\text{Sp}_{C_0(X)}(f) = f(X) \cup \{0\} \quad \forall f \in C_0(X).$$

# Spectral Radius

## Definition

Let  $x \in A$ . The spectral radius of  $x$  is

$$\rho(x) = \sup\{|\lambda|; \lambda \in \text{Sp}_A(x)\}.$$

## Remark

We always have  $\rho(x) \leq \|x\|$ .

## Proposition (Gel'fand-Mazur)

For all  $x \in A$ ,

$$\rho(x) = \lim_{n \rightarrow \infty} \sqrt[n]{\|x^n\|}.$$

## Proposition

Let  $A$  be a  $C^*$ -algebra. If  $x \in A$  is normal (i.e.,  $x^*x = xx^*$ ), then

$$\rho(x) = \|x\|.$$

## Proof.

- As  $A$  is a  $C^*$ -algebra and  $x^*x = xx^*$ , we have

$$\|x^2\| = \|(x^2)^*x^2\|^{\frac{1}{2}} = \|(x^*x)^*(x^*x)\|^{\frac{1}{2}} = \|x^*x\| = \|x\|^2.$$

- An induction shows that  $\|x^{2^n}\| = \|x\|^{2^n} \forall n \in \mathbb{N}$ .
- By Gel'fand-Mazur's result,

$$\rho(x) = \lim_{n \rightarrow \infty} \|x^{2^n}\|^{\frac{1}{2^n}} = \|x\|.$$



# Spectral Radius

## Remark

If  $A$  is a Banach algebra with an antilinear involution  $x \rightarrow x^*$ , we call  $C^*$ -norm any norm such that

$$\|x\|^2 = \|x^*\|^2 = \|x^*x\| \quad \forall x \in A.$$

## Proposition

*If  $A$  is a  $C^*$ -algebra, then its norm is its unique  $C^*$ -norm.*

## Proof.

- If  $x \in A$ , for any  $C^*$ -norm we have

$$\|x\| = \sqrt{\|x^*x\|} = \sqrt{\rho(x^*x)}.$$

- The spectral radius  $\sqrt{\rho(x^*x)}$  does not depend on the norm, so this uniquely defines the  $C^*$ -norm. □

## Corollary

*Every  $*$ -isomorphism between  $C^*$ -algebras is isometric.*

## Proof.

- Let  $\phi : A_1 \rightarrow A_2$  be a  $*$ -isomorphism between  $C^*$ -algebras  $A_1$  and  $A_2$ .
- In this case  $x \mapsto \|\phi(x)\|_{A_2}$  is a  $C^*$ -norm on  $A_1$ .
- Therefore, it agrees with the original  $C^*$ -norm of  $A_1$ .
- That is,  $\phi$  is isometric.



## Setup

- $A$  is a Banach algebra with unit  $1$  such that  $\|1\| = 1$ .
- $x \in A$  and we set  $S = \text{Sp}_A(x)$  (this a compact subset of  $\mathbb{C}$ ).
- $\Omega \subset \mathbb{C}$  is an open containing  $S$ .
- $\text{Hol}(\Omega)$  is the algebra of holomorphic functions on  $\Omega$ , equipped with the topology of uniform convergence on compact sets.



# Holomorphic Functional Calculus

## Remarks

- By Cauchy's formula, if  $f \in \text{Hol}(\Omega)$ , then

$$f(z) = \frac{1}{2i\pi} \int_{\Gamma} f(\lambda)(\lambda - z)^{-1} d\lambda \quad \forall z \in S,$$

Here  $\Gamma$  is any oriented contour in  $\Omega$  whose interior contains  $S$ .

- The map  $\lambda \rightarrow (\lambda - z)^{-1}$  is analytic on  $\mathbb{C} \setminus S$ .

## Definition

If  $f \in \text{Hol}(\Omega)$ , then we define

$$f(x) := \frac{1}{2i\pi} \int_{\Gamma} f(\lambda)(\lambda - x)^{-1} d\lambda \in A,$$

where the integral is meant as a Riemann integral.

## Remark

The integral does not depend on  $\Gamma$ .

## Theorem (Holomorphic Functional Calculus)

- 1 The map  $f \rightarrow f(x)$  is a continuous unital homomorphism of algebras from  $\text{Hol}(\Omega)$  to  $A$ .
- 2 If  $f$  and  $g$  are elements of  $\text{Hol}(\Omega)$  that agree on  $S$ , then  $f(x) = g(x)$ .
- 3 For all  $f \in \text{Hol}(\Omega)$ , we have

$$\text{Sp}_A f(x) = f(S).$$

# Example

## Example

Let  $f(z) = \sum_{n \geq 0} a_n z^n$  have convergence radius  $R > \|x\|$ .

- Let  $\Gamma = \{|\lambda| = r\}$  with  $\|x\| < r < R$ . By definition,

$$f(x) = \frac{1}{2i\pi} \int_{\Gamma} f(\lambda)(\lambda - x)^{-1} d\lambda.$$

- If  $|\lambda| > \|x\|$ , then  $\|\lambda^{-1}x\| < 1$ , and so

$$(\lambda - x)^{-1} = \lambda^{-1}(1 - \lambda^{-1}x)^{-1} = \sum_{n \geq 0} \lambda^{-(n+1)} x^n.$$

- Thus,

$$f(x) = \sum_{n \geq 0} \left( \frac{1}{2i\pi} \int_{\Gamma} \lambda^{-(n+1)} f(\lambda) d\lambda \right) x^n = \sum_{n \geq 0} a_n x^n.$$

## Proposition

Let  $B$  be a closed subalgebra of  $A$  containing the unit  $1_A$ .

- ① If  $x \in B$  is invertible in  $A$ , then  $x^{-1}$  belongs to  $B$ .
- ② For all  $x \in B$  we have

$$\text{Sp}_B(x) = \text{Sp}_A(x).$$

## Proof.

- If  $x \in B$  is invertible in  $A$ , then

$$x^{-1} = \frac{1}{2i\pi} \int_{\Gamma} \frac{\lambda^{-1}}{\lambda - x} d\lambda \in B$$

- Thus,  $x \in B^{-1}$ , and hence  $A \cap B^{-1} = B^{-1}$ .
- Therefore, if  $x \in B$ , then

$$\lambda \notin \text{Sp}_A(x) \iff x - \lambda \in A^{-1} \iff x - \lambda \in B^{-1} \iff \lambda \notin \text{Sp}_B(x).$$

That is,  $\text{Sp}_B(x) = \text{Sp}_A(x)$ .



## Proposition

Assume that  $A$  is a  $C^*$ -algebra.

- ① If  $x \in A$  is unitary (i.e.,  $x^*x = xx^* = 1$ ), then  $\text{Sp}_A(x) \subset S^1$ .
- ② If  $x \in A$  is selfadjoint (i.e.,  $x^* = x$ ), then  $\text{Sp}_A(x) \subset \mathbb{R}$ .

## Proof of (1).

Let  $x \in A$  be unitary.

- As  $\|x\|^2 = \|x^*x\| = \|1_A\| = 1$ , we have  $\text{Sp}_A(x) \subset \overline{D(0,1)}$ .
- Since  $x^{-1} = x^*$  is unitary, we also have  $\text{Sp}_A(x^{-1}) \subset \overline{D(0,1)}$ .
- If  $f(z) = z^{-1}$ , then

$$\text{Sp}_A(x) = \text{Sp}_A(f(x^{-1})) = f(\text{Sp}_A(x^{-1})) \subset \mathbb{C} \setminus D(0,1).$$

- Thus,

$$\text{Sp}_A(x) \subset \overline{D(0,1)} \cap [\mathbb{C} \setminus D(0,1)] = S^1.$$



## Proof of (2).

Let  $x \in A$  be selfadjoint.

- Set  $u = \exp(ix) = \sum \frac{1}{n!}(ix)^n$ . We have

$$u^* = \sum \frac{1}{n!}((ix)^n)^* = \sum \frac{1}{n!}(-ix^*)^n = \exp(-ix).$$

- As  $f \rightarrow f(x)$  is an algebra homomorphism and  $\exp(-iz)\exp(iz) = \exp(iz)\exp(-iz) = 1$ , we have

$$uu^* = u^*u = \exp(-ix)\exp(ix) = 1.$$

- Thus,  $u$  is unitary, and hence  $\text{Sp}_A(u) \subset \mathbb{S}^1$ .
- By spectral permanence,

$$\text{Sp}_A(u) = \text{Sp}_A(\exp(ix)) = \exp(i \text{Sp}_A(x)) \subset \mathbb{S}^1$$

- It follows that  $\text{Sp}_A(x) \subset \mathbb{R}$ .



# Gel'fand Spectrum

## Setup

$A$  is a unital commutative  $C^*$ -algebra.

## Definition

- A character of  $A$  is a linear map  $\chi : A \rightarrow \mathbb{C}$  such that

$$\begin{aligned}\chi(xy) &= \chi(x)\chi(y) & \forall x, y \in A, \\ \chi(1_A) &= 1.\end{aligned}$$

- The set of characters of  $A$  is called the Gel'fand spectrum of  $A$  and is denoted  $\text{Sp } A$ .

## Remark

It can be shown that characters are in one-to-one correspondence with maximal ideals of  $A$ .

## Example

Let  $A = C(X)$ , where  $X$  is a compact space.

- Every  $x \in X$  defines a character of  $C(X)$  by

$$\chi_x(f) = f(x), \quad f \in C(X).$$

- The map  $X \ni x \rightarrow \chi_x \in \text{Sp}(A)$  is one-to-one.
- It can be shown it is onto.
- Therefore, the characters of  $C(X)$  are in one-to-one correspondence with the points of  $X$ .



# Gel'fand Spectrum

The relationship between Gel'fand spectrum and the spectra of the points of  $A$  is provided by the following result.

## Proposition

For all  $x \in A$ , we have

$$\mathrm{Sp}_A(x) = \{\chi(x); \chi \in \mathrm{Sp}(A)\}.$$

## Proof.

Set  $S = \{\chi(x); \chi \in \mathrm{Sp}(A)\}$ .

- If  $\lambda \in \mathbb{C} \setminus \mathrm{Sp}_A(x)$  and  $\chi \in \mathrm{Sp} A$ , then

$$\chi((x - \lambda)^{-1})(\chi(x) - \lambda) = \chi((x - \lambda)^{-1}(x - \lambda)) = \chi(1) = 1.$$

- In particular,  $\chi(x) \neq \lambda$  for all  $\chi \in \mathrm{Sp}(A)$ , and so  $\lambda \notin S$ .
- Thus,  $\mathbb{C} \setminus \mathrm{Sp}_A(x) \subset \mathbb{C} \setminus S$ , i.e.,  $S \subset \mathrm{Sp}_A(x)$ .
- It can be shown that  $\mathrm{Sp}_A(x) \subset S$  (see Arveson, Thm. 1.9.5), and hence  $S = \mathrm{Sp}_A(x)$ . □

## Corollary

If  $\chi \in \text{Sp}(A)$ , then

$$\chi(x^*) = \overline{\chi(x)} \quad \forall x \in A.$$

## Proof.

Let  $x \in A$ .

- If  $x^* = x$ , then  $\chi(x) \in \text{Sp}_A(x) \subset \mathbb{R}$ .
- In general,  $x = x_1 + ix_2$ , with  $x_i^* = x_i$ . Then

$$\begin{aligned}\chi(x) &= \chi(x_1 + ix_2) = \chi(x_1) + i\chi(x_2), \\ \chi(x^*) &= \chi(x_1 - ix_2) = \chi(x_1) - i\chi(x_2).\end{aligned}$$

- As  $\chi(x_1)$  and  $\chi(x_2)$  are in  $\mathbb{R}$ , we see that  $\chi(x^*) = \overline{\chi(x)}$ . □

## Setup

- $A^*$  = topological dual of  $A$ . Banach space with norm,

$$\|\varphi\| := \sup_{\|x\|=1} |\langle \varphi, x \rangle|, \quad \varphi \in A^*.$$

- $\Omega$  = unit sphere  $A^*$ .
- By Banach-Alaoglu theorem  $\Omega$  is compact with respect to the weak-\* topology, i.e., the topology of pointwise convergence.

## Proposition

$\text{Sp } A$  is a closed subset of  $\Omega$ , and hence is compact with respect to the weak-\* topology.

## Proof.

- Let  $\chi \in A$ . If  $x \in A$ , then  $\chi(x) \in \text{Sp}_A(x)$ , and hence  $|\chi(x)| \leq \|x\|$ .
- This shows that  $\chi \in A^*$  and  $\|\chi\| \leq 1$ .
- As  $\chi(1) = 1$ , we see that  $\|\chi\| = 1$ , i.e.,  $\chi \in \Omega$ . Thus,  $\text{Sp}(A) \subset \Omega$ .
- By definition  $\text{Sp}(A)$  is the intersection of the sets,

$$\{\varphi \in A^*; \varphi(1) = 1\}, \quad \{\varphi \in A^*; \varphi(x)\varphi(y) = \varphi(xy)\}, \quad x, y \in A.$$

- There are closed subsets of  $A^*$  w.r.t. the weak-\* topology.
- Therefore,  $\text{Sp}(A)$  is closed with respect to that topology.  $\square$

# Gel'fand Transform

## Definition

The Gel'fand transform of  $A$  is the map

$$G_A : A \longrightarrow C(\operatorname{Sp}(A)), \quad x \longrightarrow \hat{x},$$

where

$$\hat{x}(\chi) = \chi(x), \quad x \in A, \quad \chi \in \operatorname{Sp}(A).$$

## Remark

$G_A$  is an algebra homomorphism.

## Theorem (Gel'fand-Naimark)

$G_A$  is an (isometric)  $*$ -isomorphism from  $A$  onto  $C(\operatorname{Sp} A)$ .

## Proof.

- Let  $x \in A$ . If  $\chi \in \text{Sp}(A)$ , then

$$\overline{\hat{x}(\chi)} = \overline{\chi(x)} = \chi(x^*) = (x^*)^\wedge.$$

This shows that  $G_A$  is a  $*$ -homomorphism.

- We have

$$\text{Sp}_A(x) = \{\chi(x); \chi \in \text{Sp}(A)\} = \hat{x}(\text{Sp}(A)) = \text{Sp}_{C(\text{Sp}_A)}(\hat{x}).$$

- In particular,  $\rho(x) = \rho(\hat{x})$  for all  $x \in A$ .
- Thus,

$$\|x\|^2 = \|x^*x\| = \rho(x^*x) = \rho(\widehat{x^*x}) = \rho(\widehat{\bar{x}}\hat{x}) = \|\widehat{\bar{x}}\hat{x}\| = \|\hat{x}\|^2.$$

It follows that  $G_A$  is isometric.

- It can be shown that  $G_A(A)$  separates the points of  $\text{Sp}(A)$ , and so  $G_A(A) = C(\text{Sp}(A))$  by Stone-Weierstrass theorem.  $\square$

# Gel'fand Transform

## Consequence

The Gel'fand transform provides a one-to-one correspondence,

$$\begin{array}{ccc} \{\text{Compact Hausdorff spaces}\} & & \{\text{Unital commutative } C^*\text{-algebras}\} \\ X & \xrightarrow{\quad} & C(X) \\ \text{Sp } A & \xleftarrow{\quad} & A \end{array}$$

## Remark

It can be shown this is actually an equivalence of categories.

## Consequence

We may regard unital  $C^*$ -algebras as the noncommutative analogue of compact spaces.

# Gel'fand Transform. Non-unital Case

## Setup

$A$  is a possibly non-unital  $\mathbb{C}^*$ -algebra.

## Definition

- A character of  $A$  is any non-zero algebra homomorphism  $\chi : A \rightarrow \mathbb{C}$ .
- The Gel'fand spectrum of  $A$  is the set of all characters of  $A$ . It is denoted  $\text{Sp}(A)$ .

## Remarks

- If  $A$  is unital and  $\chi : A \rightarrow \mathbb{C}$  is an algebra homomorphism, then  $\chi(1) = 1$ .
- If  $A^+$  is the unitalization of  $A$ , then any character  $\chi : A \rightarrow \mathbb{C}$  uniquely extends to a character  $\tilde{\chi} : A^+ \rightarrow \mathbb{C}$  by letting

$$\tilde{\chi}(x + \lambda \cdot 1) = \chi(x) + \lambda, \quad x \in A, \lambda \in \mathbb{C}.$$



# Gel'fand Transform. Non-unital Case

## Proposition

$\text{Sp}(A)$  is contained in the (closed) unit ball  $B_{A^*}$  of  $A^*$  and is locally compact w.r.t. the weak-\* topology.

## Proof.

- If  $\chi \in A$ , then  $\|\chi\| \leq \|\tilde{\chi}\| = 1$ . Thus  $\text{Sp}(A) \subset B_{A^*}$ .
- We have  $\text{Sp}(A) = K \setminus 0$ , where

$$K = \bigcap_{x,y \in A} \{\varphi \in B_{A^*}; \varphi(xy) = \varphi(x)\varphi(y)\}.$$

- $B_{A^*}$  is compact w.r.t. the weak-\* topology.
- $K$  is weak-\* closed, and hence is weak-\* compact.
- A basis of the weak \*-topology of  $\text{Sp}(A)$  is given by the compact sets,

$$\{\chi \in K; |\chi(x)| \geq \epsilon\}, \quad x \in A \setminus 0, \epsilon > 0.$$

- Thus,  $\text{Sp}(A)$  is weak-\* locally compact.



# Gel'fand Transform. Non-unital Case

## Definition

The Gel'fand transform of  $A$  is the map

$$G_A : A \longrightarrow C_0(\operatorname{Sp}(A)), \quad x \longrightarrow \hat{x},$$

where

$$\hat{x}(\chi) = \chi(x), \quad x \in A, \quad \chi \in \operatorname{Sp}(A).$$

## Remark

- If  $x \in A$ , then  $\hat{x} \in C_0(\operatorname{Sp}(A))$ , since, for any  $\epsilon > 0$ ,

$$\begin{aligned} \{\chi \in \operatorname{Sp}(A); |\hat{x}(\chi)| < \epsilon\} &= \{\chi \in \operatorname{Sp}(A); |\chi(x)| < \epsilon\} \\ &= \operatorname{Sp}(A) \setminus \{\chi \in \operatorname{Sp}(A); |\chi(x)| \geq \epsilon\}. \end{aligned}$$

- Here  $\{\chi \in \operatorname{Sp}(A); |\chi(x)| \geq \epsilon\}$  is a compact set of  $\operatorname{Sp}(A)$  (see previous slide).
- Thus,  $|\hat{x}| < \epsilon$  outside some compact set.

# Gel'fand Transform. Non-unital Case

## Theorem (Gel'fand-Naimark)

$G_A$  is an (isometric)  $*$ -isomorphism from  $A$  onto  $C_0(\operatorname{Sp} A)$ .

## Consequence

We have a one-to-one correspondence,

$$\begin{array}{ccc} \{\text{Locally compact Hausdorff spaces}\} & & \{\text{Commutative } C^*\text{-algebras}\} \\ X & \longrightarrow & C_0(X) \\ \operatorname{Sp} A & \longleftarrow & A \end{array}$$

# Continuous Functional Calculus

## Setup

- $A$  = unital  $C^*$ -algebra.
- $x \in A$  is normal, i.e.,  $x^*x = xx^*$ , and  $S = \text{Sp}_A(x)$ .
- $\mathcal{P}$  =  $*$ -algebra of polynomials  $\sum c_{mn}z^m\bar{z}^n$ .

## Definition

If  $f = \sum c_{mn}z^m\bar{z}^n \in \mathcal{P}$ , we set

$$f(x) := \sum c_{mn}x^m(x^*)^n.$$

## Facts

- $\mathcal{P} \ni f \rightarrow f(x) \in A$  is a  $*$ -homomorphism of algebras.
- Set  $\mathcal{B}$  be its range. This is sub- $*$ -algebra of  $A$ .
- Let  $B$  be the closure of  $\mathcal{B}$  in  $A$ . This is a unital sub- $C^*$ -algebra; this is the (unital)  $C^*$ -algebra generated by  $x$ .

## Remark

- $B$  is a unital commutative  $C^*$ -algebra.
- Therefore, its Gel'fand transform  $G_B : B \rightarrow C(\text{Sp}(B))$  is a  $*$ -isomorphism.

## Lemma

Set  $\xi = G_B(x)$ . Then:

- (i)  $f(x) = G_B^{-1}(f \circ \xi)$  for all  $f \in \mathcal{P}$ .
- (ii)  $\xi(\text{Sp}(B)) = S$ .

Proof of (i).

- $G_B : B \rightarrow C(\operatorname{Sp}(B))$  is a  $*$ -isomorphism.
- Therefore, if  $f = \sum c_{mn} z^m \bar{z}^n \in \mathcal{P}$ , then

$$\begin{aligned} G_B(f(x)) &= \sum c_{mn} G(x^m (x^*)^n) \\ &= \sum c_{mn} G(x)^m \overline{G(x)}^n \\ &= \sum c_{mn} \xi^m \bar{\xi}^n = f \circ \xi. \end{aligned}$$

- Thus,  $f(x) = G_B^{-1}(f \circ \xi)$ .



## Proof of (ii).

- We have

$$\xi(\operatorname{Sp}(B)) = \operatorname{Sp}_{C(\operatorname{Sp}(B))}(\xi) = \operatorname{Sp}_{C(\operatorname{Sp}(B))}(G_B(x))$$

- As  $G_B : B \rightarrow C(\operatorname{Sp}(B))$  is an isomorphism, we have

$$\operatorname{Sp}_{C(\operatorname{Sp}(B))}(G(x)) = \operatorname{Sp}_B(x)$$

- By spectral permanence,

$$\operatorname{Sp}_B(x) = \operatorname{Sp}_A(x) = S.$$

- Thus,

$$\xi(\operatorname{Sp}(B)) = S.$$



# Continuous Functional Calculus

## Definition

If  $f \in C(S)$ , we define

$$f(x) := G_B^{-1}(f \circ \xi).$$

## Remark

If  $f \in C(S)$ , then  $f \circ \xi$  is well defined, since  $\xi(\text{Sp}(B)) = S$ .

## Theorem (Continuous Functional Calculus)

- 1 The map  $\Phi : x \rightarrow f(x)$  is an isometric  $*$ -homomorphism from  $C(S)$  to  $A$ , whose image is the  $C^*$ -algebra generated by  $x$ .
- 2 If  $f = \sum c_{mn} z^m \bar{z}^n \in \mathcal{P}$ , then

$$\phi(f) = f(x) = \sum c_{mn} x^m (x^*)^n.$$

- 3 For all  $f \in C(S)$ , we have

$$\text{Sp}_A f(x) = f(S).$$



## Proof of (1)+(2).

- The 2nd part is immediate. In particular,  $\mathcal{B} \subset \Phi(C(S))$ .
- As  $G_B^{-1}$  maps onto  $B$ , we have  $\Phi(C(S)) \subset B$ .
- The map  $\Psi : C(S) \ni f \rightarrow f \circ \xi \in C(\operatorname{Sp}(B))$  is a  $*$ -homomorphism.
- As  $\xi(\operatorname{Sp}(B)) = S$ , for any  $f \in C(S)$ , we have

$$\|f \circ \xi\|_{C(\operatorname{Sp} B)} = \sup_{\chi \in \operatorname{Sp} B} |f(\xi(\chi))| = \sup_{z \in S} |f(z)| = \|f\|_{C(S)}.$$

- Thus,  $\Psi$  is an isometric  $*$ -homomorphism.
- As  $\Phi = G_B^{-1} \circ \Psi$ , it follows that  $\Phi$  is an isometric  $*$ -homomorphism.
- In particular, the range of  $\Phi$  is closed.
- As  $\mathcal{B} \subset \Phi(C(S)) \subset B$  and  $\mathcal{B}$  is dense in  $B$ , it follows that  $\Phi(C(S)) = B$ .



## Proof of (3).

Let  $f \in C(S)$ . Need to show that  $\text{Sp}_A(f(x)) = f(S)$ .

- We have  $\text{Sp}_{C(S)} f = f(S)$ .
- $\Phi$  is a  $*$ -isomorphism from  $C(S)$  onto  $B$ . Thus,

$$\text{Sp}_B(f(x)) = \text{Sp}_B(\Phi(f)) = \text{Sp}_{C(S)} f = f(S).$$

- By spectral permanence,

$$\text{Sp}_A(f(x)) = \text{Sp}_B(f(x)) = f(S).$$



## Remark

- The map  $C(S) \ni f \rightarrow f(x) \in A$  is a  $*$ -homomorphism.
- Thus, if  $f$  is real-valued, then  $f(x)$  is selfadjoint.

## Remark

- The homomorphism  $C(S) \ni f \rightarrow f(x) \in A$  is continuous.
- Thus, if  $(f_k) \subset \mathcal{P}$  converges uniformly on  $S$  to  $f$ , then

$$f(x) = \lim_{k \rightarrow \infty} f_k(x).$$

- This gives an alternative definition of  $f(x)$ .